arXiv:math/0110220v1 [math.AG] 19 Oct 2001

SMOOTH CURVES IN FAMILIES OF CALABI-YAU THREEFOLDS IN HOMOGENEOUS SPACES

ANDREAS LEOPOLD KNUTSEN

Abstract. We show the existence of geometrically rigid smooth curves of different degrees and genera in Calabi-Yau threefolds that are complete intersections in homogeneous spaces. In the special cases of Calabi-Yau complete intersections in projective spaces, our results are an improvement of previously known results.

1. Introduction

In this note we study embeddings of smooth complex projective curves into (smooth) Calabi-Yau threefolds. Such embeddings, and Calabi-Yau threefolds in general, have in the past decade been objects of extended interest in both algebraic geometry and physics. The goal of counting such curves (especially rational) has inspired the development of quantum cohomology and lead to the discovery of surprising relations between algebraic geometry and the theory of mirror symmetry.

Related problems are to prove the existence and finiteness of such curves. Clemens [Cl1] constructed isolated rational curves in general quintic threefolds of arbitrarily high degree. Katz [Ka] extended this result to all degrees and went on to deduce that for $d \leq 7$ the general quintic threefold contains only finitely many rational curves of degree d. This was extended to degrees $d \leq 9$ by Nijsse [Ni] and independetly by Johnsen and Kleiman [Jo-Kl]. Clemens has conjectured that the finiteness should hold in any degree [Cl2].

Oguiso [Og] proved that the general complete intersection Calabi-Yau threefold ($CICY$) of type $(2,3)$ in $\mathbf{P}^5$ also contains smooth isolated rational curves of any degree. This result was then extended to the other $CICY$s by Ekedahl, Johnsen and Sommervoll [E-J-S], and independently by Kley [Kl1], who also have results for isolated elliptic curves.

The most general result concerning the existence of smooth curves in $CICY$s is due to Kley [Kl2], who constructs curves of arbitrary degree and bounded genera (however the approach yields isolated smooth curves only for $g = 0$ and 1).

In this paper we improve these results (lowering the bound on the degrees for fixed genera) and extend them to 6 families of Calabi-Yau threefolds that are not complete intersections in projective space.

It is well known that there are five types of Calabi-Yau complete intersection threefolds ($CICY$s) in projective space, namely the intersection types (5) in $\mathbf{P}^4$, $(3,3)$ and $(4,2)$ in $\mathbf{P}^5$, $(3,2,2)$ in $\mathbf{P}^6$ and $(2,2,2,2)$ in $\mathbf{P}^7$.

We will pay special attention to families of Calabi-Yau threefolds in $\mathbf{P}^\mu$, for $6 \leq \mu \leq 10$, which are complete intersections in certain homogeneous spaces. We now briefly fix some notation and refer to [Mu1] and [Mu2] for further details.

We use the following convention: For a vector spave V^i of dimension i, we write $G(r, V^i)$ (resp. $G(V^i, r)$) for the Grassmann variety of r-dimensional subspaces (resp. quotient spaces) of V.

1991 *Mathematics Subject Classification.* 14J28.

Key words and phrases. curves, $K3$ surfaces, Calabi-Yau threefolds.

The variety $\Sigma^{10}_{12} \subseteq \mathbf{P}^{15}$ is a 10-dimensional spinor variety of degree 12. Let V^{10} be a 10-dimensional vector space with a nondegenerate second symmetric tensor λ. Then Σ^{10}_{12} is one of the two components of the subset of $G(V^{10}, 5)$ consisting of 5-dimensional totally isotropic quotient spaces [1].

The variety $\Sigma^{6}_{16} \subseteq \mathbf{P}^{13}$ is the Grassmann variety of 3-dimensional totally isotropic quotient spaces of a 6-dimensional vector space V^6 with a nondegenerate second skew-symmetric tensor σ. It has dimension 6 and degree 16.

Finally, $\Sigma^{5}_{18} = G/P \subseteq \mathbf{P}^{13}$, where G is the automorphism group of the Cayley algebra over $\mathbf{C}$ and P is a maximal parabolic subgroup. The variety has dimension 5 and degree 18.

By adjunction, we get the following families of Calabi-Yau threefolds in $\mathbf{P}^r$ as complete intersections in homogeneous spaces (we use the notation $(a_1, ..., a_n) \cap \Sigma$ for a complete intersection of type $(a_1, ..., a_n)$ in Σ):

r	intersection type of Calabi-Yau threefold
7	$(3,1,1) \cap G(2, V^5)$
8	$(2,2,1) \cap G(2, V^5)$
9	$(2, 1^6) \cap \Sigma^{10}_{12}$
10	$(2, 1^4) \cap G(V^6, 2)$
11	$(2,1,1) \cap \Sigma^{6}_{16}$
12	$(2,1) \cap \Sigma^{5}_{18}$

We will prove the existence of geometrically rigid smooth curves of different degrees and genera on each of these Calabi-Yau threefolds. We are also able to improve the results of Kley in [Kl2] for $CICY$s. The following is the complete result we will prove in this paper:

Theorem 1.1. *Let $d \geq 1$ and $g \geq 0$ be integers. Then in any of the following cases the general Calabi-Yau threefold X of a particular type contains a geometrically rigid smooth curve of degree d and genus g:*

(a) $X = (5) \subseteq \mathbf{P}^4$: $d^2 \geq 12g - 3$ *for* $0 \leq g \leq 12$, *and* $d > g/2 + 6$ *for* $13 \leq g \leq 34$.
(b) $X = (4,2) \subseteq \mathbf{P}^5$: $(d,g) = (3,1)$ *and* $(5,2)$, $d^2 \geq 16g$ *for* $0 \leq g \leq 15$, *and* $d > g/2 + 8$ *for* $16 \leq g \leq 30$.
(c) $X = (3,3) \subseteq \mathbf{P}^5$: $(d,g) = (3,1)$ *and* $(5,2)$, $d^2 \geq 16g$ *for* $0 \leq g \leq 15$, *and* $d > g/2 + 8$ *for* $16 \leq g \leq 30$.
(d) $X = (3,2,2) \subseteq \mathbf{P}^6$: $(d,g) = (3,1)$ *and* $(5,2)$, $d^2 \geq 16g$ *for* $0 \leq g \leq 4$, *and* $d > g + 4$ *for* $5 \leq g \leq 14$.
(e) $X = (2,2,2,2) \subseteq \mathbf{P}^7$: $d^2 \geq 16g$ *for* $0 \leq g \leq 3$, *and* $d > g + 4$ *for* $4 \leq g \leq 6$.
(f) $X = (3,1,1) \cap G(2, V^5) \subseteq \mathbf{P}^7$: $d^2 \geq 20g - 4$ *for* $0 \leq g \leq 4$, *and* $d > g + 5$ *for* $5 \leq g \leq 18$.
(g) $X = (2,2,1) \cap G(2, V^5) \subseteq \mathbf{P}^8$: $d^2 \geq 20g - 4$ *for* $0 \leq g \leq 4$, *and* $d > g + 5$ *for* $5 \leq g \leq 8$.
(h) $X = (2, 1^6) \cap \Sigma^{10}_{12} \subseteq \mathbf{P}^9$: $d^2 > 24g$ *for* $0 \leq g \leq 5$, *and* $d > g + 6$ *for* $6 \leq g \leq 10$.
(i) $X = (2, 1^4) \cap G(V^6, 2) \subseteq \mathbf{P}^{10}$: $d^2 \geq 28g - 3$ *for* $0 \leq g \leq 6$, *and* $d > g + 7$ *for* $7 \leq g \leq 12$.
(j) $X = (2,1,1) \cap \Sigma^{6}_{16} \subseteq \mathbf{P}^{11}$: $d^2 \geq 32g$ *for* $0 \leq g \leq 7$, *and* $d > g + 8$ *for* $8 \leq g \leq 14$.
(k) $X = (2,1) \cap \Sigma^{5}_{18} \subseteq \mathbf{P}^{12}$: $d^2 \geq 36g$ *for* $0 \leq g \leq 8$, *and* $d > g + 9$ *for* $9 \leq g \leq 16$.

---

[1] A quotient $f : V \to V'$ is totally isotropic with respect to λ if $(f \otimes f)(\lambda)$ is zero on $V' \otimes V'$.

Remark 1.2. In the cases above with $g = 0$ and 1, the same result is true for infinitesimally rigid smooth curves. For $g = 0$ this is an immediate consequence of equation (9) below. For $g = 1$, it follows from the approach in [Kl1, Sect. 3.2].

We will use the same technique that has been used in [Kl1], [Kl2], [Ka], [Og] and [E-J-S] for the $CICY$s, namely we start with a $K3$ surface of a suitable intersection type in one of these homogeneous spaces containing a particular smooth curve and construct a (nodal) Calabi-Yau threefold of a corresponding intersection type in the same ambient space containing the surface. Then we argue that this implies that the generic smooth Calabi-Yau threefold of the same intersection type contains a smooth geometrically rigid smooth curve of the same degree and genus.

1.1. Conventions and definitions.

The ground field is the field of complex numbers. We say a curve C in a variety V is *geometrically rigid* in V if the space of embedded deformations of C in V is zero-dimensional. If furthermore this space is reduced, we say that C is *infinitesimally rigid* or *isolated* in V.

A *curve* will always be reduced and irreducible.

By a $K3$ surface is meant a reduced and irreducible surface S with trivial canonical bundle and such that $H^1(\mathcal{O}_S) = 0$. In particular $h^2(\mathcal{O}_S) = 1$ and $\chi(\mathcal{O}_S) = 2$.

Line bundles and divisors on a $K3$ surface are used with little or no distinction, as well as the multiplicative and additive notation. Linear equivalence of divisors is denoted by $\sim$.

1.2. Acknowledgements.

The author thanks T. Johnsen for asking the question, and H. Clemens and H. P. Kley for useful discussions. Thanks are also due to H. Murakami, who was kind enough to translate relevant passages of [Mu2] from Japanese to English to the author, and to S. Mukai, who eventually provided the author with a preliminary English translation.

A part of this work was done during the author's stay at the Mittag-Leffler Institute in Djursholm, Sweden, and the last details were settled during the author's stay at the Department of Mathematics at the University of Utah, Salt Lake City, USA. The author wishes to thank both institutions for their hospitality and warm atmosphere.

The author has been supported by a grant from The Research Council of Norway.

2. BN GENERAL $K3$ SURFACES

By the results in [Mu1], a generic projective $K3$ surface in $\mathbf{P}^\mu$, for $\mu = 6$, 7, 8, 9 and 10, is a complete intersection in one of the homogeneous spaces described above.

The question whether a projective $K3$ surface is contained in such a homogeneous space as a complete intersection is related to the existence of a certain rigid rank two bundle on the surface. In the cases $\mu = 6$, 8, 9 and 10, this vector bundle can be obtained by using special divisors on a curve.

By a *polarized* $K3$ surface is meant a pair (S, L) where S is a $K3$ surface and L is a base point free line bundle on S.

It turns out that the existence of this vector bundle depends on the line bundle L that gives the polarization of the surface. We recall the following definition of Mukai:

Definition 2.1. [Mu2] *A polarized $K3$ surface (S, L) of genus μ is said to be* Brill-Noether (BN) general *if the inequality $h^0(M)h^0(N) < h^0(L) = \mu + 1$ holds for any pair (M, N) of non-trivial line bundles such that $M \otimes N \simeq L$.*

Clearly the polarized $K3$ surfaces which are BN general form a 19-dimensional Zariski open subset in the moduli space of polarized $K3$ surfaces of a fixed genus μ.

The following theorem is due to Mukai:

Theorem 2.2. [Mu2] *The projective models of BN general polarized $K3$ surfaces of small genus are as follows:*

genus	*projective model of BN general polarized $K3$ surface*
2	$S_2 \longrightarrow \mathbf{P}^2$ *double covering with branch sextic*
3	$(4) \subseteq \mathbf{P}^3$
4	$(2,3) \subseteq \mathbf{P}^4$
5	$(2,2,2) \subseteq \mathbf{P}^5$
6	$(1,1,1,2) \cap G(2,V^5) \subseteq \mathbf{P}^6$
7	$(1^8) \cap \Sigma_{12}^{10} \subseteq \mathbf{P}^7$
8	$(1^6) \cap G(V^6,2) \subseteq \mathbf{P}^8$
9	$(1^4) \cap \Sigma_{16}^{6} \subseteq \mathbf{P}^9$
10	$(1^3) \cap \Sigma_{18}^{5} \subseteq \mathbf{P}^{10}$

In the next section we will study the possible combinations of pairs (d,g) for a smooth curve C of degree d and genus g on a BN general $K3$ surface. We will obtain results as in [Kn].

3. SMOOTH CURVES ON BN GENERAL $K3$ SURFACES

We first recall the main result in [Kn]:

Theorem 3.1. [Kn, Thm. 1.1] *Let $n \geq 2$, $d > 0$, $g \geq 0$ be integers. Then there exists a $K3$ surface S of degree $2n$ in $\mathbf{P}^{n+1}$ containing a smooth curve C of degree d and genus g if and only if*

- (i) $g = d^2/4n + 1$ *and there exist integers $k, m \geq 1$ and $(k,m) \neq (2,1)$ such that $n = k^2 m$ and $2n$ divides kd,*
- (ii) $d^2/4n < g < d^2/4n + 1$ *except in the following cases*
  - (a) $d \equiv \pm 1, \pm 2 \pmod{2n}$,
  - (b) $d^2 - 4n(g-1) = 1$ *and* $d \equiv n \pm 1 \pmod{2n}$,
  - (c) $d^2 - 4n(g-1) = n$ *and* $d \equiv n \pmod{2n}$,
  - (d) $d^2 - 4n(g-1) = 1$ *and* $d-1$ *or* $d+1$ *divides* $2n$,
- (iii) $g = d^2/4n$ *and d is not divisible by $2n$,*
- (iv) $g < d^2/4n$ *and* $(d,g) \neq (2n+1, n+1)$.

Furthermore, in case (i) S can be chosen such that $\operatorname{Pic} S = \mathbf{Z}\frac{2n}{dk}C = \mathbf{Z}\frac{1}{k}H$ *and in cases (ii)-(iv) such that* $\operatorname{Pic} S = \mathbf{Z}H \oplus \mathbf{Z}C$, *where H is the hyperplane section of S.*

If $n \geq 4$, S can be chosen to be scheme-theoretically an intersection of quadrics in cases (i), (iii) and (iv), and also in case (ii), except when $d^2 - 4n(g-1) = 1$ *and* $3d \equiv \pm 3 \pmod{2n}$ *or* $d^2 - 4n(g-1) = 9$ *and* $d \equiv \pm 3 \pmod{2n}$, *in which case S has to be an intersection of both quadrics and cubics.*

Specializing to the three cases of complete intersections in projective space, we get the result in Theorem 9.1, in the same paper, which we include in Theorem 3.12 below.

For all possible triples of integers (n,d,g), with $n = 5$, 6, 7, 8 and 9, $d > 0$ and $g \geq 0$, we will now investigate which triples are also possible for smooth curves on BN general $K3$ surfaces. This means that we have to check the conditions in Definition 2.1.

We have an easy result for the cases $d^2 - 4n(g-1) = 0$.

Lemma 3.2. *Let S be as in Theorem 3.1(i). Then S is BN general if and only if $2n$ divides d (equivalently $k = 1$).*

Proof. Clearly if S is BN general, H cannot be divisible, so we must have $k = 1$.

Conversely, if $k = 1$, then $\operatorname{Pic} S = \mathbf{Z}H$, so S is clearly BN general. ☐

The surfaces in Theorem 3.1 with $d^2 - 4n(g-1) > 0$ can all be chosen with the property that $\operatorname{Pic} S = \mathbf{Z}H \oplus \mathbf{Z}C$. So we can restrict ourselves to studying the BN generality of such surfaces. We therefore make the following definition.

Definition 3.3. *For any integers $n \geq 2$, $d > 0$ and $g \geq 0$ satisfying $g < d^2/4n + 1$, the surface $S(n,d,g)$ will denote a polarized $K3$ surface with polarization H (that is H is base point free), such that $H^2 = 2n$ and*

$$\operatorname{Pic} S = \mathbf{Z}H \oplus \mathbf{Z}C,$$

where C is a divisor satisfying $C.H = d$ and $C^2 = 2(g-1)$.

We will make use of the following:

Lemma 3.4. *Let H and C be as in the definition of $S(n,d,g)$ above.*

(a) [Kn] *H is ample unless $d^2 - 4n(g-1) = n$ and $d \equiv n \pmod{2n}$, or $d^2 - 4n(g-1) = 4n$ and $d \equiv 0 \pmod{2n}$.*

(b) [Kn, Prop. 4.4] *C is nef unless $g = 0$ or $(d,g) = (2n+1, n+1)$.*

(c) [Og, Thm. 3],[Kn, Prop. 4.4] *If $g = 0$, or if we are not in one of the two following cases:*

(i) $(d,g) = (2n+1, n+1)$,

(ii) $d^2 - 4n(g-1) = 1$ *and* $d-1$ *or* $d+1$ *divides* $2n$,

then the generic member of $|C|$ is a smooth curve.

Clearly, for our purposes, we could directly assume that H is very ample and that C is base point free, but we will however work with the more general case $S(n,d,g)$ above.

We want to find necessary and sufficient conditions for $S(n,d,g)$ to be BN general.

Assume $H \sim D + F$ for $D > 0$ and $F > 0$. Then we can write $D = aH - bC$ and $F = (1-a)H + bC$, for some integers a, b. Since D and F are effective, one easily sees that we can assume that

$$a \geq 1 \ \text{ and } b \geq 1.$$

The first result is the following:

Lemma 3.5. *If $d > n + g$, then either $h^0(D)$ or $h^0(F)$ is zero.*

Proof. If $b \geq a$, then $h^0(D) = h^0(aH - bC) \leq h^0(bH - bC)$. If $b < a$, then $h^0(F) = h^0(bC - (a-1)H) \leq h^0((a-1)C - (a-1)H)$. So it suffices to show that $h^0(aH - aC) = 0$ for all $a \in \mathbf{Z} - \{0\}$.

Assume that $B \in |aH - aC|$ for some $a \in \mathbf{Z} - \{0\}$. Since $B^2 = 2a^2(n - d + g - 1) < -2a^2 < 0$, the basis divisor of $|B|$ must have in its support a smooth, rational curve Γ such that $\Gamma.B < 0$. Write

$$B \sim m\Gamma + R, \quad R \geq 0, \quad 1 \leq m \leq B.H = a(2n - d),$$

with Γ not appearing as a fixed component of R.

If $R = 0$, then $B = m\Gamma \sim a(H - C)$. Since $B^2 < -2a^2$, we must have $m > a$, but this would mean that $H - C$ is divisible, which is impossible, since $H - C$ is part of a basis for $\operatorname{Pic} S$.

So $R > 0$ and $m < B.H$. We also have $B.\Gamma \geq -2m$ and $\Gamma.H < B.H/m$, since H is ample by Lemma 3.4. Define

$$B' := B + (B.\Gamma)\Gamma,$$

Then ${B'}^2 = B^2$ and

$$-B.H < B'.H = B.H + (B.\Gamma)(\Gamma.H) < B.H,$$

so $(B'.H)^2 < (B.H)^2$.

Setting $B_0 := H - C$ and $B'_0 := B_0 + (B_0.\Gamma)\Gamma$, we get

$$B \sim aB_0 \quad \text{and} \quad B' \sim aB'_0,$$

and also ${B'_0}^2 = {B_0}^2$ and $(B'_0.H)^2 < (B_0.H)^2$.

By the Hodge index theorem we have

$$(B'_0.H)^2 - {B'_0}^2 H^2 \geq 0,$$

with equality occurring if and only if

$$(B'_0.H)H \sim 2nB'_0.$$

But the latter is impossible, since ${B'_0}^2 = {B_0}^2 < -2$.

So we have

$$0 \neq (B'_0.H)^2 - 2n{B'_0}^2 = |\operatorname{disc}(H, B'_0)| < (B_0.H)^2 - 2n{B_0}^2 = |\operatorname{disc}(H, B_0)|,$$

which is impossible, since $\operatorname{Pic} S \simeq \mathbf{Z}H \oplus \mathbf{Z}B_0$.

So $|aH - aC| = \emptyset$. □

Corollary 3.6. *$S = S(n, d, g)$ is BN general for $d > n + g$.*

When $g = 0$, we also have an easy result:

Lemma 3.7. *$S(n, d, 0)$ is BN general.*

Proof. Letting D and F be as above, we clearly have $h^0(F) = h^0((1-a)H + bC) = 0$ unless $a = 1$, in which case $h^0(F) = h^0(bC) = 1$. We then have $h^0(D) = h^0(H - bC) < h^0(H)$, since $|H|$ is without fixed components, and we get $h^0(M)h^0(N) < h^0(H)$ □

When $g = 1$, more interesting things start to happen. It is already known, for instance, that a complete intersection $K3$ surface of type $(2, 2, 2)$ in $\mathbf{P}^5$ cannot contain a smooth elliptic curve of degree 3, whereas there are indeed $K3$ surfaces in $\mathbf{P}^5$ containing such elliptic curves (see e.g. [SD] and [Jo-Kn]).

So let $g = 1$ and D and F be as above. Since all members of $|bC|$ are of the form $C_1 + ... + C_b$, with each $C_i \in |C|$ (see [SD, Prop. 2.6]), we have $h^0(F) = h^0((1-a)H + bC) = 0$ unless $a = 1$, in which case $h^0(F) = h^0(bC) = b + 1$ (again by [SD, Prop. 2.6]). We now want to compute $h^0(D) = h^0(H - bC)$. We will show that

$$h^0(D) = \begin{cases} \frac{1}{2}D^2 + 2 & \text{if } D^2 \geq -2, \\ 0 & \text{if } D^2 < -2. \end{cases} \tag{1}$$

By standard theory on linear systems on $K3$ surfaces and Riemann-Roch, this is an immediate consequence of the following:

Lemma 3.8. (a) *If $D^2 \geq 0$ then D is effective and nef.*

(b) *If $D^2 = -2$, then $|D|$ contains as its only element a smooth rational curve.*

(c) *If $D^2 < -2$, then $|D| = \emptyset$.*

Proof. We first show that D is effective when $D^2 = 2(n - bd) \geq -2$. By Riemann-Roch, either D or $-D$ is effective. Since $n \geq bd - 1$, we get $H.D = 2n - bd > 0$, whence D must be effective.

If either $D^2 \geq 0$ and D is not nef, or $D^2 = -2$ and $|D|$ contains no reduced, irreducible elements, or $D^2 < -2$ and $D > 0$, there has to exist a smooth, rational curve Γ such that $\Gamma.D < 0$. As above, we can write

$$D \sim m\Gamma + R, \quad m \geq 1, \quad R \geq 0,$$

with Γ not appearing as a fixed component of R. Furthermore, since $\operatorname{Pic} S \simeq \mathbf{Z}D \oplus \mathbf{Z}C$, we must have $R > 0$.

Grant for the time being the following

Claim: In all cases, $0 < \Gamma.C < D.C/m$.

By this we have

$$0 < |\operatorname{disc}(C, \Gamma)| = (\Gamma.C)^2 < (D.C)^2 = |\operatorname{disc}(C, D)|,$$

and this is impossible, since $\operatorname{Pic} S \simeq \mathbf{Z}D \oplus \mathbf{Z}C$. We have reached a contradiction, which shows that the statements (a), (b) and (c) are true. ☐

Proof of claim. Since $|C|$ contains a smooth elliptic curve, we must have $\Gamma.C \geq 0$. If $\Gamma.C = 0$, write $\Gamma \sim xH + yC$, and get $xd = 0$, so $x = 0$ and $\Gamma \sim yC$, which is impossible, since $\Gamma^2 = -2$.

Now $\Gamma.C = \frac{1}{m}(D.C - R.C)$, so it suffices to show that $R.C \neq 0$.

Assume, to get a contradiction, that $R.C = 0$. Then $D.C = m\Gamma.C = (H - bC).C = d$. Writing $\Gamma \sim xH + yC$, we easily find from $\Gamma^2 = -2$ and $\Gamma.C = d/m$, that

$$m = x = 1 \quad \text{and} \quad y = -\frac{n+1}{d}$$

so that

$$\Gamma \sim H - \frac{n+1}{d}C, \qquad d \mid n+1.$$

We calculate $\Gamma.D = n - 1 - bd$, and since $\Gamma.D < 0$ by hypothesis, we have

$$n < bd + 1. \tag{2}$$

We also calculate $R.H = (\frac{n+1}{d} - b)C.H = n + 1 - bd$. By Lemma 3.4, if H is not ample, then $d = 1$, but this would imply that H is not base point free, by standard criteria for base point freeness of line bundles on a $K3$ surface (see e.g. [Kn, Lemma 2.4]). Since R is effective and H is ample, we must have

$$n > bd - 1. \tag{3}$$

Combining (2) and (3) we get $n = bd$, but then d cannot divide $n + 1$, a contradiction. This concludes the proof of the claim. ☐

We now have, using $D^2 = 2(n - bd)$ and (1),

$$h^0(D)h^0(F) = \begin{cases} (n - bd + 2)(b + 1) & \text{if } bd \leq n + 1, \\ 0 & \text{if } bd > n + 1, \end{cases} \tag{4}$$

This gives us

Lemma 3.9. *If $g = 1$ and $n \geq 2$, then $S = S(n, d, 1)$ is BN general for $n \leq 9$ if and only if $d \geq 3$ and $n = 2$ or 3, $d \geq 4$ and $n = 4$ or 5, $d \geq 5$ and $n = 6$ or 7, $d \geq 6$ and $n = 8$ or 9.*

We will now consider curves of higher genera.

By the two following lemmas, we can restrict to considering the cases $d \leq n$.

Lemma 3.10. *Let $n \geq 2, d \geq 1, g \geq 0$ be integers satisfying $g < d^2/4n + 1$. Then there are integers $d_0 \geq 1$ and $g_0 \geq 0$ such that $d_0 \equiv d \pmod{2n}$ and $d_0^2 - 4n(g_0 - 1) = d^2 - 4n(g-1)$, and satisfying one of the two following conditions:*

(a) *$g_0 < d_0 - n$,*
(b) *$d_0 \leq 2n$, $d_0 - n \leq g_0 < d_0^2/4n + 1$.*

Proof. If $d \leq 2n$, we are done. If $d \geq 2n+1$, let $d' := d - 2n$ and $g' = g - d + n$. Then, by the assumptions on d and g, we have $d' \geq 1$, $g' \geq 0$, $d' \equiv d \pmod{2n}$ and $d'^2 - 4n(g'-1) = d^2 - 4n(g-1)$.

If $g' < d' - n$, we are in case (a) with $(d_0, g_0) = (d', g')$, and there cannot exist any integers as in (b), for setting $d'' = d' - 2n$ and $d''^2 - 4n(g''-1) = d^2 - 4n(g-1)$, we get $g'' = g' - d' + n < 0$.

If $g' \geq d' - n$, we continue the procedure until we either find a pair (d_0, g_0) as in (a) or (b). □

Lemma 3.11. *Let $n \geq 2, d \geq 1, g \geq 0$ be integers satisfying $g < d^2/4n + 1$.*

(a) Let d_0 and g_0 be as in Lemma 3.10. Then $S(n,d,g)$ is BN general if and only if $S(n, d_0, g_0)$ is BN general.

(b) Assume $n-d+g \geq 0$. Then $S(n,d,g)$ is BN general if and only if $S(n, 2n-d, n-d+g)$ is BN general.

Proof. Let $d = d_0 + 2nk$, for an integer $k \geq 0$. Then (i) follows from a change of basis $C \mapsto C - kH$.

Similarly, (ii) follows from a change of basis $C \mapsto H - C$. □

By (a) we can assume that $d \leq 2n$ and that $d \leq n + g$ (the case $d > n + g$ being already treated by Lemma 3.5).

By (b) we can in addition assume that $d \leq n$.

By the Lemmas 3.7 and 3.9, this means that we have already treated the cases $d = n + g$ and $d = n + g - 1$ for $2 \leq n \leq 9$.

Now we require H to be very ample, that is we require n, d and g to be as in Theorem 3.1, except that we a priori allow the case (ii) (d) to happen (since this contradicts the base point freeness of C and not the very ampleness of H, as is seen in [Kn, Prop. 4.4]). But one easily checks that the case (ii) (d) does not occur for $n \leq 9$, when H is very ample. So the only cases that are left to check for $n \leq 9$ are the following:

$$\begin{array}{rcllll} (n,d,g) & = & (6,6,2), & & & \\ & & (7,7,2), & (7,6,2), & (7,8,3), & \\ & & (8,8,2), & (8,7,2), & (8,6,2), & (8,8,3), \\ & & (9,9,2), & (9,8,2), & (9,7,2), & (9,9,3). \end{array}$$

By computing the product [2]

$$h^0(H-C)h^0(C) = (\frac{1}{2}(H-C)^2 + 2)(\frac{1}{2}C^2 + 2) = (n - d + g + 1)(g+1),$$

---

[2] Here we use the fact that $h^1(C) = h^1(H-C) = 0$, which follows from the fact that both $|C|$ and $|H-C|$ contain an irreducible member, which one can check using arguments as in the proof of Lemma 3.5.

one finds that this is greater than or equal to $h^0(H) = n+2$ in all these cases, except

$$(n, d, g) = (8, 8, 2) \ \text{ and } \ (9, 9, 2).$$

We now show that in these two cases $S(n, d, g)$ is in fact BN general.

As usual, write $D := aH - bC$ and $F := (1-a)H + bC$, for two integers a, $b \geq 1$, and assume $D, F > 0$. Since H is (very) ample, we have

$$D.H = 2na - bd > 0$$

and

$$F.H = (1-a)2n + bd > 0,$$

which gives

$$\frac{2n}{d}(a-1) < b < \frac{2n}{d}a. \tag{5}$$

Since C is nef, we have

$$D.C = da - 2(g-1)b \geq 0$$

and

$$F.C = d(1-a) + 2(g-1)b \geq 0,$$

which gives

$$\frac{2b}{d}(g-1) \leq a \leq \frac{2b}{d}(g-1) + 1. \tag{6}$$

For $(n, d, g) = (8, 8, 2)$ and $(9, 9, 2)$, (5) gives

$$2a - 2 < b < 2a,$$

so $b = 2a - 1$, and the right hand inequality of (6) gives $a < 2$ in both cases, so we must have $a = b = 1$, and we only need to check the product $h^0(C)h^0(H-C)$ as above.

We summarize these results, and get the following result for the existence of smooth curves on BN general $K3$ surfaces of genus $\mu = 6, ..., 10$. For completeness we also include the cases with $\mu = 3$, proved by Mori in [Mo], and the cases $\mu = 4$ and 5, proved in [Kn].

Theorem 3.12. *Let $d > 0$ and $g \geq 0$ be integers. Then:*

(i) *(Mori* [Mo]*) There exists a smooth quartic surface X in $\mathbf{P}^3$ containing a smooth curve C of degree d and genus g if and only if (a) $g = d^2/8 + 1$, or (b) $g < d^2/8$ and $(d, g) \neq (5, 3)$. Furthermore, (a) holds if and only if $\mathcal{O}_X(1)$ and $\mathcal{O}_X(C)$ are dependent in* $\operatorname{Pic} X$*, in which case C is a complete intersection of X and a hypersurface of degree $d/4$.*

(ii) *There exists a $K3$ surface X of type $(2, 3)$ in $\mathbf{P}^4$ containing a smooth curve C of degree d and genus g if and only if (a) $g = d^2/12+1$, (b) $g = d^2/12+1/4$ or (c) $g < d^2/12$ and $(d, g) \neq (7, 4)$. Furthermore, (a) holds if and only if $\mathcal{O}_X(1)$ and $\mathcal{O}_X(C)$ are dependent in* $\operatorname{Pic} X$*, in which case C is a complete intersection of X and a hypersurface of degree $d/6$.*

(iii) *There exists a $K3$ surface X of type $(2, 2, 2)$ in $\mathbf{P}^5$ containing a smooth curve C of degree d and genus g if and only if (a) $g = d^2/16 + 1$ and d is divisible by 8, (b) $g = d^2/16$ and $d \equiv 4 \pmod 8$, or (c) $g < d^2/16$ and $(d, g) \neq (9, 5)$. Furthermore, (a) holds if and only if $\mathcal{O}_X(1)$ and $\mathcal{O}_X(C)$ are dependent in* $\operatorname{Pic} X$*, in which case C is a complete intersection of X and a hypersurface of degree $d/8$.*

(iv) *There exists a* $K3$ *surface* X *of type* $(1,1,1,2) \cap G(2, V^5)$ *in* $\mathbf{P}^6$ *containing a smooth curve* C *of degree* d *and genus* g *if and only if (a)* $g = d^2/20 + 1$*, (b)* $g = d^2/20 + 1/5$*, or (c)* $g < d^2/20$ *and* $(d, g) \neq (11, 6)$*. Furthermore, (a) holds if and only if* $\mathcal{O}_X(1)$ *and* $\mathcal{O}_X(C)$ *are dependent in* $\operatorname{Pic} X$*, in which case* C *is a complete intersection of* X *and a hypersurface of degree* $d/10$*.*

(v) *There exists a* $K3$ *surface* X *of type* $(1^8) \cap \Sigma^{10}_{12}$ *in* $\mathbf{P}^7$ *containing a smooth curve* C *of degree* d *and genus* g *if and only if (a)* $g = d^2/24 + 1$*, or (b)* $g < d^2/24$ *and* $(d, g) \neq (13, 7)$*. Furthermore, (a) holds if and only if* $\mathcal{O}_X(1)$ *and* $\mathcal{O}_X(C)$ *are dependent in* $\operatorname{Pic} X$*, in which case* C *is a complete intersection of* X *and a hypersurface of degree* $d/12$*.*

(vi) *There exists a* $K3$ *surface* X *of type* $(1^6) \cap G(V^6, 2)$ *in* $\mathbf{P}^8$ *containing a smooth curve* C *of degree* d *and genus* g *if and only if (a)* $g = d^2/28 + 1$*, (b)* $g = d^2/28 + 3/28$*, or (c)* $g < d^2/28$ *and* $(d, g) \neq (15, 8)$*. Furthermore, (a) holds if and only if* $\mathcal{O}_X(1)$ *and* $\mathcal{O}_X(C)$ *are dependent in* $\operatorname{Pic} X$*, in which case* C *is a complete intersection of* X *and a hypersurface of degree* $d/14$*.*

(vii) *There exists a* $K3$ *surface* X *of type* $(1^4) \cap \Sigma^{6}_{16}$ *in* $\mathbf{P}^9$ *containing a smooth curve* C *of degree* d *and genus* g *if and only if (a)* $g = d^2/32 + 1$ *and* d *is divisible by* 16*, (b)* $g = d^2/28$ *and* $d \equiv 8 \pmod{16}$*, or (c)* $g < d^2/32$ *and* $(d, g) \neq (17, 9)$*. Furthermore, (a) holds if and only if* $\mathcal{O}_X(1)$ *and* $\mathcal{O}_X(C)$ *are dependent in* $\operatorname{Pic} X$*, in which case* C *is a complete intersection of* X *and a hypersurface of degree* $d/16$*.*

(viii) *There exists a* $K3$ *surface* X *of type* $(1^3) \cap \Sigma^{5}_{18}$ *in* $\mathbf{P}^{10}$ *containing a smooth curve* C *of degree* d *and genus* g *if and only if (a)* $g = d^2/36 + 1$ *and* d *is divisible by* 18*, (b)* $g = d^2/36$ *and* $d \equiv \pm 6 \pmod{18}$*, or (c)* $g < d^2/36$ *and* $(d, g) \neq (19, 10)$*. Furthermore, (a) holds if and only if* $\mathcal{O}_X(1)$ *and* $\mathcal{O}_X(C)$ *are dependent in* $\operatorname{Pic} X$*, in which case* C *is a complete intersection of* X *and a hypersurface of degree* $d/18$*.*

We are now ready to prove the main theorem of this paper.

4. PROOF OF THEOREM 1.1

We recall the main setting and results of [Kl2].

Let P be a smooth projective variety, $\mathcal{E}$ a locally free sheaf of rank $\dim P - 3$ on P, and

$$s_0 \in H^0(P, \mathcal{E})$$

a regular section. Set

$$X_0 = Z(s_0).$$

Let $S \subseteq X_0$ be a regular surface (i.e. $H^1(S, \mathcal{O}_S) = 0$) and L a line bundle on S.

We make the following assumptions:

(A1) X_0 is K-trivial

(A2) The only singularities of X_0 which lie in S are m ordinary double points, and these are distinct from any singularities of S and base points of $|L|$. Furthermore

$$m \geq \dim |L| + 2.$$

(A3) $H^1(C', \mathcal{N}_{C'/P}) = 0$ for all $C' \in |L|$.

(A4) $H^0(C', \mathcal{N}_{C'/S}) \simeq H^0(C', \mathcal{N}_{C'/X_0})$ for all $C' \in |L|$.

(A5) There exists an $s \in H^0(P, \mathcal{E})$ such that $X_t := Z(s_0 + ts)$ is a smoothing of at least one of the ordinary double points of X_0.

Then one has:

Theorem 4.1. [Kl2, Thm. 1.1] *Under the assumptions (A1)-(A5) the members of* $|L|$ *deform to a length* $\binom{m-2}{\dim|L|}$ *scheme of curves which are geometrically rigid in the general deformation* $X_t = Z(s_0 + ts)$ *of* X_0*. In particular,* X_t *contains a geometrically rigid curve which is a deformation of a curve in* $|L|$

By Theorem 3.12 the possible degrees and genera of smooth curves on BN general $K3$ surfaces of low genus are classified.

For the cases where X is a $CICY$, the results in Theorem 1.1 are slight improvements of the results of Kley in [Kl2, Thm. 2.1], by using [Kn, Prop. 1.2] (which is an improvement of [Kl2, Lemma 2.3]). For the cases where X is a complete intersection in some homogeneous space different than projective space, we will have to argue that the conditions (A1)-(A5) are still satisfied. Our argument will also be valid for the $CICY$ cases.

Given a $K3$ surface S in $\mathbf{P}^\mu$ of intersection type $(a_1, ..., a_{k(\mu)})$ in some smooth ambient space P (like some projective space or homogeneous space as described above), we can construct Calabi-Yau threefolds X_0 of intersection type $(b_1, ..., b_{k(\mu)-1})$ in the same ambient space, whenever $1 \leq a_i \leq b_i$, for $i = 1, ..., k(\mu) - 1$, as follows. Choose generators g_i of degrees a_i for the ideal of S. So $S = V(g_1, \dots, g_{k(\mu)})$. For general $\alpha_{ij} \in H^0(P, \mathcal{O}_P(b_i - a_j)$ define

$$f_i := \sum \alpha_{ij} g_j$$

and

$$X_0 := V(f_1, \dots, f_{k(\mu)})$$

(here we follow [Kl1, Section 3]). If the coefficient forms are chosen in a sufficiently general way, X_0 has only m ordinary double points. The numbers m in the $CICY$ cases are given in [Kl2, Table p.6], which we here reproduce:

(b_i)	(a_j)	m	(b_i)	(a_j)	m
$(5) \subseteq \mathbf{P}^4$	$(4,1) \subseteq \mathbf{P}^4$	16	$(3,3) \subseteq \mathbf{P}^5$	$(3,2,1) \subseteq \mathbf{P}^5$	12
$(5) \subseteq \mathbf{P}^4$	$(3,2) \subseteq \mathbf{P}^4$	36	$(3,3) \subseteq \mathbf{P}^5$	$(2,2,2) \subseteq \mathbf{P}^5$	32
$(4,2) \subseteq \mathbf{P}^5$	$(4,1,1) \subseteq \mathbf{P}^5$	4	$(3,2,2) \subseteq \mathbf{P}^6$	$(3,2,1,1) \subseteq \mathbf{P}^6$	6
$(4,2) \subseteq \mathbf{P}^5$	$(3,2,1) \subseteq \mathbf{P}^5$	18	$(3,2,2) \subseteq \mathbf{P}^6$	$(2,2,2,1) \subseteq \mathbf{P}^6$	16
$(4,2) \subseteq \mathbf{P}^5$	$(2,2,2) \subseteq \mathbf{P}^5$	32	$(2,2,2,2) \subseteq \mathbf{P}^7$	$(2,2,2,1,1) \subseteq \mathbf{P}^7$	8

In the remaining cases, we have the following table:

(b_i)	(a_j)	m
$(3,1,1) \cap G(2,V^5) \subseteq \mathbf{P}^7$	$(2,1,1,1) \cap G(2,V^5) \subseteq \mathbf{P}^6$	20
$(2,2,1) \cap G(2,V^5) \subseteq \mathbf{P}^8$	$(2,1,1,1) \cap G(2,V^5) \subseteq \mathbf{P}^6$	10
$(2,1^6) \cap \Sigma^{10}_{12} \subseteq \mathbf{P}^9$	$(1^8) \cap \Sigma^{10}_{12} \subseteq \mathbf{P}^7$	12
$(2,1^4) \cap G(V^6,2) \subseteq \mathbf{P}^{10}$	$(1^6) \cap G(V^6,2) \subseteq \mathbf{P}^8$	14
$(2,1,1) \cap \Sigma^6_{16} \subseteq \mathbf{P}^{11}$	$(1^4) \cap \Sigma^6_{16} \subseteq \mathbf{P}^9$	16
$(2,1) \cap \Sigma^5_{18} \subseteq \mathbf{P}^{12}$	$(1^3) \cap \Sigma^5_{18} \subseteq \mathbf{P}^{10}$	18

Therefore, in all the cases given by Theorem 3.12, we have inclusions

$$C \subseteq S \subseteq X_0 \subseteq P,$$

where C is a smooth curve of degree d and genus g, S is a smooth projective $K3$ surface, X_0 is a nodal Calabi-Yau threefold of the corresponding type, and P is the homogeneous variety in question.

Now the conditions (A1) and (A5) are clearly satisfied. Set $L := \mathcal{O}_S(C)$. The condition (A2) is satisfied whenever

$$m \geq \dim |L| + 2 = g + 2.$$

This yields the upper bounds on the genus in the different cases of Theorem 1.1.

It remains to check the conditions (A3) and (A4).

Consider the standard short exact sequence

$$0 \longrightarrow \mathcal{T}_S \longrightarrow \mathcal{T}_P \otimes \mathcal{O}_S \longrightarrow \mathcal{N}_{S/P} \longrightarrow 0. \tag{7}$$

The key observation is the following result, which is an extension of [Kl1, Prop. 1.7] to include the cases where the ambient space P is not a projective space:

Proposition 4.2. *Let S be a BN general $K3$ surface in $\mathbf{P}^{\mu}$, $3 \leq \mu \leq 10$, and P its corresponding ambient space where it is a complete intersection. The sequence*

$$H^0(S, \mathcal{N}_{S/P}) \xrightarrow{\delta} H^1(S, \mathcal{T}_S) \xrightarrow{c} H^2(S, \mathcal{O}_S),$$

where δ arises from (7) and c is the Yoneda pairing $\times c_1(\mathcal{O}_S(1))$, is exact.

Proof. Since $\omega_S \simeq \mathcal{O}_S$, the Yoneda pairing

$$H^1(S, \mathcal{T}_S) \times H^1(S, \Omega^1_S) \longrightarrow H^2(S, \mathcal{O}_S)$$

is non-degenerate by Serre duality.

Now $H^1(S, \mathcal{T}_S)$ is the moduli space of all (analytic) deformations of S, which is 20-dimensional, and $H^0(S, \mathcal{N}_{S/P})$ is the moduli space of all deformations of S in P, whose image by δ in $H^1(S, \mathcal{T}_S)$ is 19-dimensional, since S is BN general.

It is then clear that $\operatorname{im} \delta = \ker c$. $\square$

For each $C' \in |L|$ there is an exact sequence

$$0 \longrightarrow \mathcal{N}_{C'/S} \longrightarrow \mathcal{N}_{C'/P} \longrightarrow \mathcal{N}_{S/P} \otimes \mathcal{O}_{C'} \longrightarrow 0. \tag{8}$$

Now one can check that with the help of Proposition 4.2, we can prove the following analogue to [Kl1, Lemma 1.10]:

Lemma 4.3. *Assume that $\mathcal{O}_S(1)$ and L are independent in $\operatorname{Pic} S$. Then for all $C' \in |L|$ the composition*

$$\phi : H^0(S, \mathcal{N}_{S/P}) \longrightarrow H^0(C', \mathcal{N}_{S/P} \otimes \mathcal{O}_{C'}) \longrightarrow H^1(C', \mathcal{N}_{C'/S})$$

of the restriction morphism with the connecting homomorphism arising from (8) is surjective. Furthermore, $\ker \phi$ is independent of $C' \in |L|$.

Since S is a complete intersection of type $(a_1, ..., a_k)$ in P, we have $\mathcal{N}_{S/P} \otimes \mathcal{O}_{C'} \simeq \oplus \mathcal{O}_{C'}(a_i)$. Lemma 4.3 together with (8) gives

$$H^1(C', \mathcal{N}_{C'/P}) \simeq H^1(C', \mathcal{N}_{S/P} \otimes \mathcal{O}_{C'}) \simeq \oplus H^1(C', \mathcal{O}_{C'}(a_i)),$$

In particular

$$H^1(\mathcal{N}_{S/P} \otimes \mathcal{O}_{C'}) = 0 \text{ if and only if } H^1(C', \mathcal{O}_{C'}(\min\{a_i\})) = 0.$$

By looking at the tables on the previous page, we see that we have $\min\{a_i\} = 1$ in most cases, and $\min\{a_i\} = 2$ in three of the cases where P is projective space.

By [Kn, Prop 1.2], we now get that

$$(A3) \text{ is satisfied if and only if } d \leq 2(\mu - 1) \text{ or } d > \mu + g - 1,$$

(where μ is the genus of S) in the cases where $\min\{a_i\} = 1$, and

$$(A3) \text{ is satisfied if and only if } d \leq 4(\mu - 1) \quad \text{or} \quad 2d > 4\mu + g - 4,$$

in the cases where $\min\{a_i\} = 2$. This gives the constraints on d in Theorem 1.1.

We now only have left to show that condition (A4) is satisfied, which means that we must show that

$$h^0(\mathcal{N}_{C'/X_0}) = h^0(\mathcal{N}_{C'/S}) = g \tag{9}$$

for all $C' \in |C|$.

This follows the lines of the proof of [Kl1, Thm. 3.5], with the help of Lemma 4.3 again. This concludes the proof of Theorem 1.1.

References

[Cl1] H. Clemens, *Homological equivalence, modulo algebraic equivalence, is not finitely generated*, Publ. Math. IHES **58**, 19-38 (1983).

[Cl2] H. Clemens, *Some Results on Abel-Jacobi Mappings*, in *Topics in transcendental algebraic geometry*, Princeton Univ. Press, Princeton, NJ (1984).

[E-J-S] T. Ekedahl, T. Johnsen, D. E. Sommervoll, *Isolated rational curves on K3-fibered Calabi-Yau threefolds*, Manuscr. Math. **99**, no.1, 111-133 (1999).

[Ka] S. Katz, *On the finiteness of rational curves on quintic threefolds*, Comp. Math. **60**, 151-162 (1986).

[Jo-Kl] T. Johnsen, S. Kleiman, *Rational curves of degree at most* 9 *on a general quintic threefold*, Comm. in Algebra **24** (8), 2721-2753 (1996).

[Jo-Kn] T. Johnsen, A. L. Knutsen, *$K3$ projective models in scrolls*, Max-Planck-Institut für Mathematik Preprint Series 2001.

[Kn] A. L. Knutsen, *Smooth curves on projective $K3$ surfaces*, to appear in Math. Scand. (preprint available at math.AG/9805140).

[Kl1] H. P. Kley, *Rigid curves in complete intersection Calabi-Yau threefolds*, Comp. Math. **123**, no. 2, 185-208 (2000).

[Kl2] H. P. Kley, *On the existence of curves in K-trivial threefolds*, Preprint, alg-geom 9811099 (1998).

[Mo] S. Mori, *On degrees and genera of curves on smooth quartic surfaces in* $\mathbf{P}^3$, Nagoya Math. J. **96**, 127-132 (1984).

[Mu1] S. Mukai, *Curves, $K3$ surfaces and Fano 3-folds of genus ≤ 10*, in *Algebraic Geometry and Commutative Algebra in Honor of Masayoshi NAGATA*, 357-377 (1987).

[Mu2] S. Mukai, *New development of theory of Fano 3-folds: vector bundle method and moduli problem*, Sugaku **47**, 125-144 (1995).

[Ni] P. G. J. Nijsse, *Clemens' conjecture for octic and nonic curves*, Indag. Math. (N.S.) **6**, no. 2, 213-221 (1995).

[Og] K. Oguiso, *Two remarks on Calabi-Yau Moishezon threefolds*, J. für die Reine und Angew. Math. **452** (1994), 153-162.

[SD] B. Saint-Donat, *Projective models of $K3$ surfaces*, Amer. J. Math. **96**, 602-639 (1974).

DEPT. OF MATHEMATICS, UNIVERSITY OF BERGEN, JOHS. BRUNSGT 12, 5008 BERGEN, NORWAY
E-mail address: `andreask@mi.uib.no`